

Craniotomy and Resection of Intrinsic Lesion

	intrinsic Lesion	
Facility:		

(Affix patient identification label here)						
URN:						
Family Name:						
Given Names:						
Address:						
Date of Birth:	Sex: M	F				

A. Interpreter / cultural needs	
An Interpreter Service is required?	☐ Yes ☐ No
If yes, is a qualified Interpreter present?	☐ Yes ☐ No
A Cultural Support Person is required?	☐ Yes ☐ No
If yes, is a Cultural Support Person present?	Yes No
R Condition and treatment	

The doctor has explained that you have the following condition: (Doctor to document in patient's own words)

This condition requires the following procedure. (Doctor to document - include site and/or side where relevant to the procedure)

A Craniotomy and resection of Intrinsic Lesion is performed to remove a lesion from within the brain.

C. Risks of this procedure

There are risks and complications with this procedure. They include but are not limited to the following.

Common risks and complications include:

- Infection, requiring antibiotics and further treatment.
- Minor pain, bruising and/or infection from IV cannula site. This may require treatment with antibiotics.
- Bleeding can occur and may require a return to the operating room. Bleeding is more common if you have been taking blood thinning drugs such as Warfarin, Asprin, Clopidogrel (Plavix or Iscover) or Dipyridamole (Persantin or Asasantin).
- · Heart attack due to the strain on the heart.
- Stroke or stroke like complications may occur causing neurological deficits such as weakness in the face, arms and legs. This could be temporary or permanent.
- Injury to the brain, important nerves or blood vessels. This can lead to stroke like complications.
- Memory disturbance or confusion. This could be temporary or permanent.
- Fluid leakage from around the brain may occur through the wound after the operation. This may require further surgery.

- Epilepsy which may require medication. This condition may be temporary or permanent.
- Abnormal sensations such as pins and needles, numbness or pain may occur from the wound after the operation. This may be temporary or permanent.
- All of the lesion may not be removed by surgery and may require further treatment such as radiotherapy and/or chemotherapy.

Uncommon risks and complications include:

- Skull deformity and/or poor cosmetic result may occur requiring further surgery at a later stage.
- Decrease in the normal body salt concentration.
 This may require admission to intensive care and further treatment.
- Small areas of the lung may collapse, increasing the risk of chest infection. This may need antibiotics and physiotherapy.
- Increase risk in obese people of wound infection, chest infection, heart and lung complications, and thrombosis.
- Blood clot in the leg (DVT) causing pain and swelling. In rare cases part of the clot may break off and go to the lungs.

Rare risks and complications include:

- Meningitis may occur requiring further treatment and antibiotics.
- Visual disturbance which may be temporary or permanent.
- Coma.
- Death as a result of this procedure is possible.

D.	Significant	risks	and	procedure
	options			

(Doctor to document in space provided. Continue in Medical Record if necessary.)
E. Risks of not having this procedure
(Doctor to document in space provided. Continue in Medical Record if necessary.)



Craniotomy and Resection of

Intrinsic Lesion Facility:

	(Affix patient ide	ntification labe	l here)			
URN:						
Family Name:						
Given Names:						
Address:						
Date of Birth:			Sex:	М	F	

F. Anaesthetic
This procedure may require an anaesthetic. (Doctor to document type of anaesthetic discussed)
G. Patient consent

I acknowledge that the doctor has explained;

- my medical condition and the proposed procedure, including additional treatment if the doctor finds something unexpected. I understand the risks, including the risks that are specific to me.
- the anaesthetic required for this procedure. I understand the risks, including the risks that are specific to me.
- other relevant procedure options and their associated risks.
- my prognosis and the risks of not having the procedure.
- that no guarantee has been made that the procedure will improve my condition even though it has been carried out with due professional care.
- the procedure may include a blood transfusion.
- tissues and blood may be removed and could be used for diagnosis or management of my condition, stored and disposed of sensitively by the hospital.
- if immediate life-threatening events happen during the procedure, they will be treated accordingly.
- a doctor other than the Specialist Neurosurgeon may conduct the procedure. I understand this could be a doctor undergoing further training.

I have been given the following Patient

nformation Sheet/s;
☐ About your Anaesthetic
☐ Craniotomy & Resection of Intrinsic Lesion
Luca able to sak questions and raise concerns

- I was able to ask questions and raise concerns with the doctor about my condition, the proposed procedure and its risks, and my treatment options. My questions and concerns have been discussed and answered to my satisfaction.
- I understand I have the right to change my mind at any time before the procedure, including after I have signed this form but, preferably following a discussion with my doctor.

On the basis of the above statements.

I request to have the procedure Name of Patient/ Substitute decision maker and relationship:
Signature:
Date:
Substitute Decision-Maker: Under the Powers of Attorney Act 1998 and/or the Guardianship and Administration Act 2000. If the patient is an adult and unable to give consent, an authorised decision-maker must give consent on the patient's behalf.

I have explained to the patient all the above points under the Patient Consent section (G) and I am of the opinion that the patient/substitute decision-

H. Doctor's statement

maker has understood the information.
Name of
Doctor:
Designation:
Designation.
Ciamatuma
Signature:
Date:
Name of
Anaesthetist:
Designation:
Designation.
Ciamatuma
Signature:
Date:

I. Interpreter's statement
I have given a sight translation in
(state the patient's language here) of the consent form and assisted in the provision of any verbal and written information given to the patient/parent or guardian/substitute decision-maker by the doctor. Name of
Interpreter:
Signature:
Date:



Consent Information - Patient Copy Craniotomy and Resection of Intrinsic Lesion

1. What is a Craniotomy and Resection of Intrinsic Lesion?

A Craniotomy and resection of Intrinsic Lesion is performed to remove a lesion from within the brain.

A cut is made over the area of the lesion. A segment of bone will be removed. If the lesion is not seen on the surface of the brain, a cut is made into the brain to expose the lesion. A computerised navigation system maybe used to locate the lesion.

The lesion is removed. The skull bone is put back and is closed with metal plates and screws.

The cut is closed with stitches or staples.





Medical illustration Copyright © Nucleus Medical Art. All Rights Reserved.

2. My anaesthetic

This procedure will require a General Anaesthetic.

See **About your Anaesthetic** information sheet for information about the anaesthetic and the risks involved. If you have any concerns, talk these over with your doctor.

If you have not been given an information sheet, please ask for one.

3. What are the risks of this specific procedure?

There are risks and complications with this procedure. They include but are not limited to the following.

Common risks and complications include:

- Infection, requiring antibiotics and further treatment.
- Minor pain, bruising and/or infection from IV cannula site. This may require treatment with antibiotics.
- Bleeding can occur and may require a return to the operating room. Bleeding is more common if you have been taking blood thinning drugs such as Warfarin, Asprin, Clopidogrel (Plavix or Iscover) or Dipyridamole (Persantin or Asasantin).

- · Heart attack due to the strain on the heart.
- Stroke or stroke like complications may occur causing neurological deficits such as weakness in the face, arms and legs. This could be temporary or permanent.
- Injury to the brain, important nerves or blood vessels. This can lead to stroke like complications.
- Memory disturbance or confusion. This could be temporary or permanent.
- Fluid leakage from around the brain may occur through the wound after the operation. This may require further surgery.
- Epilepsy which may require medication. This condition may be temporary or permanent.
- Abnormal sensations such as pins and needles, numbness or pain may occur from the wound after the operation. This may be temporary or permanent.
- All of the lesion may not be removed by surgery and may require further treatment such as radiotherapy and/or chemotherapy.

Uncommon risks and complications include:

- Skull deformity and/or poor cosmetic result may occur requiring further surgery at a later stage.
- Decrease in the normal body salt concentration.
 This may require admission to intensive care and further treatment.
- Small areas of the lung may collapse, increasing the risk of chest infection. This may need antibiotics and physiotherapy.
- Increase risk in obese people of wound infection, chest infection, heart and lung complications, and thrombosis.
- Blood clot in the leg (DVT) causing pain and swelling. In rare cases part of the clot may break off and go to the lungs.

Rare risks and complications include:

- Meningitis may occur requiring further treatment and antibiotics.
- Visual disturbance which may be temporary or permanent.
- Coma.
- Death as a result of this procedure is possible.

Notes to talk to my doctor about					